



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### Corrigendum to "GLOWORM-PARA

**Citation for published version:**

Rose Vineer, H, Verschave, SH, Claerebout, E, Vercruysse, J, Shaw, DJ, Charlier, J & Morgan, ER 2021, 'Corrigendum to "GLOWORM-PARA: a flexible framework to simulate the population dynamics of the parasitic phase of gastrointestinal nematodes infecting grazing livestock" [Int. J. Parasitol. 50 (2020) 133-144]', *International Journal For Parasitology*. <https://doi.org/10.1016/j.ijpara.2021.05.001>

**Digital Object Identifier (DOI):**

[10.1016/j.ijpara.2021.05.001](https://doi.org/10.1016/j.ijpara.2021.05.001)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Publisher's PDF, also known as Version of record

**Published In:**

International Journal For Parasitology

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.





## Corrigendum

# Corrigendum to “GLOWORM-PARA: a flexible framework to simulate the population dynamics of the parasitic phase of gastrointestinal nematodes infecting grazing livestock” [Int. J. Parasitol. 50 (2020) 133–144]

H. Rose Vineer<sup>a,b,c,\*</sup>, S.H. Verschave<sup>d,e</sup>, E. Claerebout<sup>d</sup>, J. Vercruyssen<sup>d</sup>, D.J. Shaw<sup>f</sup>, J. Charlier<sup>g</sup>, E.R. Morgan<sup>a,b,h</sup>

<sup>a</sup> Veterinary Parasitology and Ecology Group, Bristol Veterinary School, University of Bristol, BS8 1TQ, UK

<sup>b</sup> Cabot Institute, Royal Fort House, University of Bristol, BS8 1UJ, UK

<sup>c</sup> Department of Infection Biology, Institute of Infection and Global Health, University of Liverpool, Leahurst Campus, Neston, Cheshire CH64 7TE, UK

<sup>d</sup> Department of Virology, Parasitology and Immunology, Faculty of Veterinary Medicine, Ghent University, Salisburylaan 133, 9820 Merelbeke, Belgium

<sup>e</sup> Department of Molecular and Cellular Biology, Harvard University, 52 Oxford Street, Cambridge, MA 02138, USA

<sup>f</sup> The Royal (Dick) School of Veterinary Studies and The Roslin Institute, University of Edinburgh, Easter Bush Campus, Roslin EH25 9RG, UK

<sup>g</sup> Kreavet, Hendrik Mertensstraat 17, 9150 Kruibeke, Belgium

<sup>h</sup> Institute for Global Food Security, Queen's University Belfast, BT9 7BL, UK

The authors regret an error in the R code used for model validation using the *Cooperia oncophora* parameters. Correcting this error results in minor changes to the validation statistics and plots for *C. oncophora*. However, conclusions remain unchanged and readers

wishing to replicate the analyses or view corrected statistics can refer the updated [Table 3](#), [Fig. 3](#), [Fig. 4](#) and [supplementary data](#) here.

The authors would like to apologise for any inconvenience caused.

**Table 3**

Validation of simulations for faecal egg counts (FEC) of *O. ostertagi* and *C. oncophora* using parasitological data of first season grazing animals on seven commercial dairy herds in Belgium.

Dataset	<i>Ostertagia ostertagi</i>				<i>Cooperia oncophora</i>			
	Error (residual sum of squares)	Linear regression	$R^2$ ( $R^2$ adjusted)	Slope (95% CI)	Error (residual sum of squares)	Linear regression	$R^2$ ( $R^2$ adjusted)	Slope (95% CI)
Herd 1	37.69	$F_{1,5} = 6.89, p=0.047$	0.58 (0.50)	0.71 (0.19–1.24)	68.17	$F_{1,5} = 1.004, p=0.362$	0.17 (–0.0007)	0.26 (0.25–0.78)
Herd 2	10.11	$F_{1,3} = 13.35, p=0.035$	0.82 (0.76)	0.14 (0.07–0.22)	68.34	$F_{1,3} = 47.11, p=0.006$	0.94 (0.92)	0.49 (0.35–0.63)
Herd 3	0.61	$F_{1,2} = 766.7, p=0.001$	1 (1)	0.71 (0.66–0.76)	6.094	$F_{1,2} = 257.4, p=0.004$	0.99 (0.99)	4.08 (–3.59–4.57)
Herd 4	8.38	$F_{1,6} = 7.00, p=0.038$	0.54 (0.46)	0.77 (0.20–1.33)	5.77	$F_{1,6} = 6.53, p = 0.043$	0.52 (0.44)	1.28 (–0.30–2.27)
Herd 5	47.31	$F_{1,3} = 15.1, p=0.030$	0.83 (0.78)	1.71 (0.85–2.57)	300.9	$F_{1,3} = 8.71, p = 0.060$	0.74 (0.66)	11.98 (–4.02–19.93)
Herd 6	9.42	$F_{1,4} = 30.35, p=0.005$	0.88 (0.85)	2.54 (1.64–3.45)	85.29	$F_{1,4} = 6.0, p=0.071$	0.60 (0.50)	6.88 (–1.37–12.39)
Herd 7	7.39	$F_{1,2} = 3.84, p = 0.189$	0.66 (0.49)	0.19 (–0.0002–0.39)	2.14	$F_{1,2} = 5.62, p = 0.141$	0.74 (0.61)	0.06 (0.01–0.11)


DOI of original article: <https://doi.org/10.1016/j.ijpara.2019.11.005>

\* Corresponding author at: Department of Infection Biology, Institute of Infection and Global Health, University of Liverpool, Leahurst Campus, Neston, Cheshire CH64 7TE, UK.


E-mail address: [hannah.vineer@liverpool.ac.uk](mailto:hannah.vineer@liverpool.ac.uk) (H. Rose Vineer).

<https://doi.org/10.1016/j.ijpara.2021.05.001>

0020-7519/© 2021 Australian Society for Parasitology. Published by Elsevier Ltd. All rights reserved.



**Fig. 3.** Observed and predicted faecal egg counts for *Cooperia oncophora* in first season grazing animals of dairy herds in Belgium. FECs were monitored for the entire length of the first grazing season. Further information on the background of this data can be found in Supplementary Table S1. Points and error bars show the observed number of eggs per gram of faeces and the corresponding 95% confidence interval obtained by bootstrapping (10,000 repeats). The dashed black line depicts the predicted FEC for a group of hosts based on a deterministic model simulation. The solid grey lines depict predictions obtained from 50 model simulations representing individual hosts, in which stochastic L3 intake and between-host variability in immune response were incorporated.



**Fig. 4.** The observed and simulated faecal egg counts (points) with 95% confidence intervals (observed = horizontal, simulated = vertical). The diagonal black line indicates hypothetical perfect agreement between the observed and simulated faecal egg counts. The grey solid line indicates the predicted slope of the regression, with 95% confidence intervals shown as grey dashed lines. Note that the 95% confidence intervals for the simulated data (estimated using the stochastic simulations shown in grey in Fig. 2, Fig. 3) are narrow and may not be easily seen due to the scale of the Y-axes.

#### Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ijpara.2021.05.001>.